

Letters to the editor

THE RELATIONSHIP BETWEEN SHOPLIFTING AND BORDERLINE PERSONALITY SYMPTOMATOLOGY AMONG INTERNAL MEDICINE OUTPATIENTS

DEAR EDITOR:

The lifetime prevalence of shoplifting in the United States is around 11 percent, and may be explained in various ways, including associations with psychiatric disorders.¹ However, few studies have examined relationships between shoplifting and borderline personality disorder (BPD), a disorder characterized by inherent impulsivity. While studies in eating disorder²⁻⁴ and kleptomania⁵ samples indicate possible associations between BPD and shoplifting, a study by Saint-Martin and Chabrol⁶ did not report the same findings. In this study, we hypothesized a relationship between shoplifting and borderline personality symptomatology.

Participants were men and women being seen for nonemergent medical care at an outpatient internal medicine clinic staffed predominantly by resident providers. We excluded individuals with compromising medical (e.g., pain), intellectual (e.g., mental retardation), cognitive (e.g., dementia), or psychiatric symptoms (e.g., psychotic) that would preclude the candidate's ability to successfully complete a survey.

During clinic hours, one of the authors (Lam) approached consecutive patients in the lobby and informally assessed exclusion criteria. With potential candidates, the recruiter reviewed the focus of the project and then invited each to complete a five-page survey.

Participants were asked to place the completed surveys into sealed envelopes and then into a collection box in the lobby.

The survey consisted of three core sections: 1) a demographic query, 2) a yes/no query about shoplifting (i.e., "Have you ever shoplifted?"), and 3) two assessments for borderline personality symptomatology—the borderline personality scale of the Personality Diagnostic Questionnaire-Revised⁷ and the Self-Harm Inventory.⁸ This project was approved by an institutional review board and completion of materials was assumed to be informed consent, which was clarified on the cover page.

Of the 471 individuals approached, 417 agreed to participate (participation rate of 88.5%). Of these, 379 completed relevant study measures. These 130 men and 249 women ranged in age from 19 to 97 years (mean 50.30, standard deviation 15.44). Most respondents (88.1%) were Caucasian, followed by African-American (7.9%), other (2.1%), Hispanic (1.1%), and Asian (0.8%). All but 6.4 percent of respondents reported having at least attained a high school diploma; 14.7 percent reported a bachelor's degree, and 14.1 percent a graduate or professional degree.

Ninety six (25.3%) indicated having shoplifted. Compared to those who denied ever having shoplifted ($M=1.61$, $SD=1.95$, $n=279$), those who indicated such behavior ($M=3.44$, $SD=2.54$, $n=96$) had statistically significantly higher PDQ-4 scores, $F(1,373)=53.46$, $p<0.001$. Similarly, compared to those who denied ever having shoplifted ($M=1.33$, $SD=2.27$,

$n=276$), those who indicated such behavior ($M=5.20$, $SD=4.64$, $n=96$) had statistically significantly higher SHI scores, $F(1,370)=113.92$, $p<0.001$. In addition, a greater proportion of those who indicated having shoplifted (36.5%) exceeded the PDQ-4 cut-off score compared to the proportion of those who denied ever having shoplifted (10.8%), $\chi^2=32.94$, $p<0.001$. Similarly, a greater proportion of those who indicated having shoplifted (49.0%) exceeded the SHI cut-off score compared to the proportion of those who denied having shoplifted (9.8%), $\chi^2=68.60$, $p<0.001$.

According to findings, there is an association between shoplifting and borderline personality symptomatology, likely affecting only a portion of individuals with this personality disorder. Viewed as an impulsive behavior, shoplifting is consistent with the construct of BPD.

This study has a number of potential limitations, including the self-report nature of all data and the tendency for self-report measures of BPD to be overinclusive. However, in this consecutive and reasonably sized sample, we examined a relationship that is rarely explored and found that shoplifting is a behavior that may be associated with BPD.

REFERENCES

1. Blanco C, Grant J, Petry NM, et al. Prevalence and correlates of shoplifting in the United States: results from the National Epidemiologic Survey on Alcohol and Related Conditions (NESARC). *Am J Psychiatry*. 2008;165:905-913.
2. Selby EA, Bulik CM, Thornton L, et al. Refining behavioral dysregulation in borderline personality using a sample of

- women with anorexia nervosa. *Pers Disord Theory Res Treat.* 2010;1:250–257.
3. Nagata T, Kawarada Y, Kiriike N, Iketani T. Multi-impulsivity of Japanese patients with eating disorders: primary and secondary impulsivity. *Psychiatry Res.* 2000;94:239–250.
 4. Suzuki K, Higuchi S, Yamada K, et al. Bulimia nervosa with and without alcoholism: a comparative study in Japan. *Int J Eat Disord.* 1994;16:137–146.
 5. Grant JE. Co-occurrence of personality disorders in persons with kleptomania: a preliminary investigation. *J Am Acad Psychiatry Law.* 2004;32:395–398.
 6. Saint-Martin C, Chabrol H. Contribution of psychopathic traits to delinquent behaviors in a sample of high-school male students. *Encephale.* 2010;36:155–158.
 7. Hyler SE. *Personality Diagnostic Questionnaire-4.* New York: New York State Psychiatric Institute; 1994.
 8. Sansone RA, Wiederman MW, Sansone LA. The Self-Harm Inventory (SHI): development of a scale for identifying self-destructive behaviors and borderline personality disorder. *J Clin Psychol.* 1998;54:973–983.
- With regards,
- Randy A. Sansone, MD;**
Charlene Lam, MD, MPH; and
Michael W. Wiederman, PhD
- Dr. Sansone is a Professor in Psychiatry and Internal Medicine at Wright State University School*

of Medicine in Dayton, Ohio, and Director of Psychiatry Education at Kettering Medical Center in Kettering, Ohio. Dr. Lam is a resident in the transition-year program at Kettering Medical Center in Kettering, Ohio. Dr. Wiederman is a Professor in Psychology in the Department of Human Relations at Columbia College in Columbia, South Carolina.

Address correspondence to:

Randy A. Sansone, MD, Sycamore Primary Care Center, 2115 Leiter Road, Miamisburg, OH 45342; Phone: 937-384-6850; Fax: 937-384-6938. E-mail: Randy.sansone@khnetwork.org

LETTERS TO THE EDITOR SUBMISSIONS

Innovations in Clinical Neuroscience welcomes submissions for possible publication in the Letters section. Letters should be no more than 500 words and five references. Type with double spacing. Letters that comment on a *Innovations in Clinical Neuroscience* article will be considered if they are received within two months of the time the article was published. All letters are reviewed by the editors and are selected based upon interest, timeliness, and pertinence, as determined by the editors. There are no guarantees a letter will be published. Letters that are not intended for publication by the authors should state this in the body of the letter. Published letters are edited and may be shortened; tables and figures are included only selectively.

Send letters to: Elizabeth Klumpp, Executive Editor, *Innovations in Clinical Neuroscience*, editorial@matrixmedcom.com. Hard copy submissions are no longer accepted.

Include the following statement with the letter: “The undersigned author transfers all copyright ownership of the enclosed/attached letter to Matrix Medical Communications in the event the work is published. The undersigned author warrants that the letter is original, is not under consideration by another journal, and has not been previously published. I sign for and accept responsibility for releasing this material on behalf of any and all co-authors.”